

Products	Active Ingredient	Persistent Activity (Days)	Target Parasites														Comments	Dosage	Slaughter Withdrawal (Days)	Administration			
			Lungworms	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Lice	Mange Mites	Horn Flies					Grubs	Beef	Lactating Dairy
CYDECTIN® Injectable	Moxidectin	14-42	●						●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	21	SubQ
CYDECTIN® Pour-On	Moxidectin	28-42	●						●		●	●	●	●	●	●	●	●	●	Do not use in calves to be processed for veal.	5 ml / 110 lbs	0	Topical
DECTOMAX® Injectable	Doramectin	21-28	●						●	●	●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 100 lbs	35	SubQ/IM
DECTOMAX® Pour-On	Doramectin	21-35	●	●	●		●		●	●	●	●	●	●	●	●	●	●	●	For use in dairy replacement heifers under 20 months of age. Not for use in veal calves.	50 ml / 1100 lbs	45	Topical
Eprinex® Pour-On	Eprinomectin	21-28	●						●		●	●	●	●	●	●	●	●	●	Do not use in calves to be processed for veal.	1 ml / 110 lbs	0	Topical
Ivermectin Pour-On	Ivermectin	14-28	●						●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 22 lbs	48	Topical
Ivermectin Injectable	Ivermectin	14-28	●						●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	35	SubQ
Ivomec® Pour-On	Ivermectin	14-28	●						●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	50 ml / 1100 lbs	48	Topical
Ivomec® Injectable	Ivermectin	14-28	●						●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	35	SubQ
Ivermectin Plus Injectable	Ivermectin/ Clorsulon	14-28	●					●	●		●	●	●	●	●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	49	SubQ
LongRange® Injectable <sup>Rx only</sup>	Eprinomectin	100-150	●	●	●	●	●						●		●	●	●	●	●	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	48	SubQ
PROHIBIT®	Levamisole		●	●	●	●	●										●	●	●	Do not use in female dairy cattle over 20 months of age.	2 ml / 100 lbs	2	Drench
Panacur® 10%	Fenbendazole		●	●	●	●	●										●	●	●	Do not use in calves to be processed for veal.	2.3 ml / 100 lbs*	8	Drench
Safe-Guard® Suspension 10%	Fenbendazole		●	●	●	●	●										●	●	●	Do not use in calves to be processed for veal.	2.3 ml / 100 lbs*	8	Drench
Safe-Guard® Paste 10%	Fenbendazole		●	●	●	●	●										●	●	●	Do not use in calves to be processed for veal.	5 g / 220 lbs	8	Oral
Synanthic® 9.06%	Oxfendazole		●	●	●		●	●									●	●	●	Do not use in female dairy cattle over 20 months of age.	2.5 ml /	7	Drench
Synanthic® 22.5%	Oxfendazole		●	●	●		●	●									●	●	●	Do not use in female dairy cattle over 20 months of age.	1 ml / 110 lbs	7	Drench
Valbazen®	Albendazole		●	●	●	●	●	●	●								●	●	●	Do not use in female dairy cattle over 20 months of age.	4 ml / 100 lbs	27	Drench

\* 2.3 ml/100 lbs or 2.3 ml/50 lbs for ascarids